



Semaglutide for Weight Loss

Overview: Semaglutide is a medication prescribed to help manage weight loss and diabetes. It mimics the effects of glucagon-like peptide-1 (GLP-1), which increases insulin production, impacts blood sugar, and supports healthy metabolism. Semaglutide has been approved by the FDA under various brand names for the treatment of chronic weight management and obesity.

Suggested Dosage:

0.5mg to 2mg of Semaglutide, injected weekly as a subcutaneous injection.

Semaglutide must be titrated or started at a lower dose and gradually increased over time, to help the body adjust to the medication. The maximum dose for Semaglutide is a 2mg subcutaneous injection weekly, though patients may require smaller doses depending on their needs, lab results, and response to the medication.

Titration schedule for 2.5mg/mL formula(s):

1. Start with a 0.25mg (0.1mL or 10 units on an insulin syringe) subcutaneous injection weekly for 4 weeks.
2. After that, increase to a 0.5mg (0.2mL or 20 units on an insulin syringe) subcutaneous injection weekly.
3. Patients may continue to increase to 1mg (0.4mL or 40 units on an insulin syringe) weekly for at least 4 weeks, and then may increase to 2mg (0.8mL or 80 units on an insulin syringe) weekly for at least 4 weeks.

Titration schedule for 1mg/mL + 0.5mg/mL Cyanocobalamin formula:

1. Start with a 0.25mg (0.25mL or 25 units on an insulin syringe) subcutaneous injection weekly for 4 weeks.
2. After that, increase to a 0.5mg (0.5mL or 50 units on an insulin syringe) subcutaneous injection weekly.
3. Patients may continue to increase to 1mg (1mL or 100 units on an insulin syringe) weekly for at least 4 weeks, and then may increase to 2mg (2mL or 200 units on an insulin syringe) weekly for at least 4 weeks.

Titration schedule for 5mg/mL + 0.5mg/mL Cyanocobalamin formula:

1. Start with a 0.25mg (0.05mL or 5 units on an insulin syringe) subcutaneous injection weekly for 4 weeks.
2. After that, increase to a 0.5mg (0.1mL or 10 units on an insulin syringe) subcutaneous injection weekly.
3. Patients may continue to increase to 1mg (0.2mL or 20 units on an insulin syringe) weekly for at least 4 weeks, and then may increase to 2mg (0.4mL or 40 units on an insulin syringe) weekly for at least 4 weeks.

A provider will guide your titration schedule and final dosage, and it's important to follow your provider's instructions. Patient protocols may vary from the general guidelines provided here.



Combination Therapies:

Semaglutide may be combined with other medications and supplements as appropriate under the supervision of a provider. The following products may be co-administered to potentiate effects:

- LIPO-C Injections
- B12 Injections
- Lipo Burn weight management products
- Bella weight management products
- LipoSculpt cream

Semaglutide Injection	Vial Size	SKU	Patient Price
2.5mg/mL	1mL	1228	\$175
2.5mg/mL	3mL	1228	\$275
1mg/mL + 0.5mg/mL Cyanocobalamin	2.5mL	1228	\$189
5mg/mL + 0.5mg/mL Cyanocobalamin	2.5mL	1228	\$289